

Confinement in the lattice Landau Gauge QCD simulation

Sadataka Furui* and Hideo Nakajima†

*School of Science and Engineering, Teikyo University, Utsunomiya 320-8551, Japan

†Department of Information science, Utsunomiya University, Utsunomiya 320-8585, Japan

Abstract. The running coupling and the Kugo-Ojima parameter of the confinement criterion are measured for the quenched SU(3) $\beta = 6.4, 6.45$, 56^4 lattice and the unquenched $\beta = 5.2$, $20^3 \times 48$ lattice of JLQCD, $\beta = 2.1$, $\kappa = 0.1357, 0.1382$, $24^3 \times 48$ lattice of CP-PACS and $\beta_{imp} = 6.76$, $am_{u,d} = 0.007, 6.83$, $am_{u,d} = 0.040$, $20^3 \times 64$ lattice of MILC collaboration.

The quenched SU(3) 56^4 lattice data suggest presence of infrared fixed point of $\alpha_s(0) = 2.5(5)$ and the approach of the ensemble of the 1st copy to the Gribov boundary. The running coupling of $q > 2\text{GeV}$ can be fitted by the perturbative QCD(pQCD) + c_1/q^2 correction. We find the Kugo-Ojima parameter $u(0) = -0.83(3)$.

The rotational symmetry of the gluon propagator of the unquenched SU(3) is partially recovered, but its magnitude depends on whether the Wilson fermion or the Kogut-Susskind(KS) fermion are coupled to the gauge field. When the sea quark mass is sufficiently light, both fermions suggest infrared fixed point $\alpha_s(0) \sim 2 - 2.5$. The Kugo-Ojima parameter of unquenched configurations with light fermion masses is consistent with $u(0) = -1.0$.

INTRODUCTION

In Landau gauge $\widetilde{MOM}$ scheme, we measure the QCD running coupling in terms of gluon dressing function $Z_A(q^2)$ and ghost dressing function $G(q^2)$, $\alpha_s(q) = \frac{g^2}{4\pi} \frac{G(\tilde{q}^2/a^2)^2 Z_A(q^2)}{\tilde{Z}_1^2}$. ($q = \sqrt{\tilde{q}^2 + \tilde{q}^4(\delta/3)}/a$, $\delta = 0$ for ordinary action and $\delta = 1$ for the improved action[5, 6].) It is a renormalization group invariant quantity, but in the finite lattice, the vertex renormalization factor $\tilde{Z}_1$ is not necessarily equal to 1 as in pQCD. We fix this value by the fit of the numerical result to the pQCD.

Colour confinement in infrared QCD is characterized by the Kugo and Ojima parameter $u(0) = -c$. The parameter c is related to the renormalization factor as $1 - c = \frac{Z_1}{Z_3} = \frac{\tilde{Z}_1}{\tilde{Z}_3}$. If the finiteness of $\tilde{Z}_1$ is proved, divergence of $\tilde{Z}_3$ is a sufficient condition. If Z_3 vanishes in the infrared, Z_1 should have higher order 0.

THE GHOST PROPAGATOR AND THE GLUON PROPAGATOR

The ghost propagator is the Fourier transform of an expectation value of the inverse Faddeev-Popov operator $\mathcal{M} = -\partial D = -\partial^2(1 - M)$

$$D_G^{ab}(x, y) = \langle \text{tr} \langle \Lambda^a x | (\mathcal{M}[U])^{-1} | \Lambda^b y \rangle \rangle \quad (1)$$

where the outmost $\langle \rangle$ denotes average over samples U .

The ghost propagators $D_G(q^2)$ of quenched and unquenched SU(3) in $\widetilde{MOM}$ scheme can be fitted by the pQCD in $q > 0.4\text{GeV}$ region, and there is no β dependence. It means that there are strong correlation between the string tension and the ghost propagator. In this presentation the gauge field of log- U type[2] is adopted. The ghost propagators are 14% larger when the U -linear definition is adopted.

The gluon propagator of 56^4 lattice is finite at 0 momentum and $Z_3(0)$ is not compatible to 0 in the present lattice size. The gluon propagator of the unquenched SU(3)[4, 5, 6] is measured by adopting the cylinder cut. Extraction of infrared physical quantities becomes difficult due to lack of symmetry of the four coordinate axes.

We observe reflection positivity violation in the unquenched gluon propagator, and in some polarization components of sample-wise quenched gluon propagator. The reflection positivity violation and the closeness of the Kugo-Ojima parameter to -1 are correlated.

THE QCD RUNNING COUPLING AND THE KUGO-OJIMA PARAMETER

The QCD running coupling $\alpha_s(q)$ of the quenched SU(3) normalized at high momentum region by the 3-loop pQCD is close to the prediction of Dyson-Schwinger calculation of $\kappa = 0.5$. The dependence of the running coupling on the sea quark mass of Wilson fermion (JLQCD, CP-PACS) and that of KS fermion(MILC) are qualitatively different. Relatively heavy Wilson fermion (JLQCD) does not show deviation from pQCD in $q > 3\text{GeV}$ region, but both fermions suggest infrared fixed point $\alpha_s(0) \sim 2 - 2.5$, when the sea quark mass is light. The left-most momentum point in the Figure 1 is to be excluded in the cone-cut due to the finite size effect.

The Kugo-Ojima parameters are summarized in Table 1. The parameter c becomes larger as the lattice size becomes large. It is about 0.83(3) in 56^4 , while in the unquenched simulation it is consistent to 1, when the fermion mass is light. In the case of U -linear definition of the gauge field, c is about 10% smaller than that of the log- U definition.

SUMMARY

We observed that the running coupling of unquenched Landau gauge QCD in the infrared is enhanced when the sea quark mass is light and the c_1/q^2 correction which appeared in the quenched simulation appears also in the unquenched simulation. The KS




FIGURE 1. The running coupling $\alpha_s(q)$ of the CP-PACS of $K_{sea} = 0.1357$ (diamonds) and that of 0.1382 (triangles). (25 samples) [left] and that of the MILC $\beta_{imp} = 6.83, am_{u,d} = 0.040$ (diamonds) and $6.76, am_{u,d} = 0.007$ (triangles). (50 samples) [right]. The DSE approach with $\alpha_0 = 2.5$ (long dashed line), pQCD (dash-dotted line) and the Orsay fit of $pQCD + c_1/q^2$ (dashed line) are also shown.

TABLE 1. The Kugo-Ojima parameter along the spacial directions c_x and that along the time axis c_t and the average c , trace divided by the dimension e/d , horizon function deviation h of quenched Wilson action (The first two rows), unquenched Wilson action (The second two rows), unquenched Wilson improved action (The third two rows), and unquenched KS improved action (The last two rows).

K_{sea} or β	c_x	c_t	c	e/d	h
$\beta = 6.4$	0.827(27)	0.827(27)	0.827(27)	0.954(1)	-0.12(3)
$\beta = 6.45$	0.809(81)	0.809(81)	0.809(81)	0.954(1)	-0.15(8)
$K_{sea} = 0.1340$	0.887(87)	0.723(38)	0.846(106)	0.930(1)	-0.084(106)
$K_{sea} = 0.1355$	1.005(217)	0.670(47)	0.921(238)	0.934(1)	-0.013(238)
$K_{sea} = 0.1357$	0.859(58)	0.763(36)	0.835(68)	0.9388(1)	-0.104(58)
$K_{sea} = 0.1382$	0.887(87)	0.723(38)	0.846(106)	0.9409(1)	-0.051(87)
$\beta_{imp} = 6.76$	1.040(111)	0.741(28)	0.965(162)	0.9325(1)	0.032(162)
$\beta_{imp} = 6.83$	0.994(141)	0.748(32)	0.932(163)	0.9339(1)	-0.002(163)

fermion of MILC collaboration (Asqtad action) has smaller sea quark mass dependence than that of Wilson fermion of CP-PACS collaboration.

This work is supported by the KEK supercomputing project No.04-106. H.N.is supported by a JSPS Grant-in-Aid for Scientific Research on Priority Area No.13135210.

REFERENCES

1. S. Furui and H.Nakajima, Phys. Rev. D**69**,074505(2004), hep-lat/0305010 and references therein.
2. S. Furui and H.Nakajima, Phys. Rev. D**70**,094504(2004), hep-lat/0403021 and references therein.
3. S. Furui and H. Nakajima, in *Confinement IV*, Ed. W. Lucha et.al., World Scientific, Singapore, p.275(2002), hep-lat/0012017.
4. S.Aoki et al., (JLQCD collaboration), Phys. Rev. D**65**,094507(2002).
5. A. AliKhan et al., (CP-PACS collaboration), Phys. Rev. D**65**,054505(2002).
6. C.W. Bernard et al., (MILC collaboration), Phys. Rev. D**64**,054506(2001).